



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE FOUR-POINT PROBLEM.

BY T. J. LOWRY, U. S. C. S., SAN FRANCISCO, CAL.

PROBLEM :—Four points in the same plane being given in position, to determine the position of any other point (or place of observation) in reference to these given points, having on the four points two angles which have no parts common except their vertices.



Let A, B, D, & C be the four given points : and let DGC & AGB be the observed angles. And then since the $\angle AEB = 2 \angle AGB$, and $AE = BE \therefore$ in $\triangle AEB$ we have BA and all the \angle s to find AE. And then in $\triangle EAC$ are known AE and AC and $\angle EAC$ to find EC and the $\angle ECA$. But

since $\angle DFC = 2 \angle DGC$, and $DF = FC$ hence in $\triangle DFC$ we have DC and all the \angle s to get FC. But the $\angle FCE = \angle ACD - \angle ACE - \angle FCD \therefore$ in $\triangle FCE$ are given FC and CE and the included angle to find FE and $\angle EFC$, then in $\triangle EGF$ all the sides are known, hence $\angle GFE$ can be found. But $\angle GFC = \angle EFC + \angle GFE \therefore$ in $\triangle GFC$ we have GF and FC and all the angles to find GC. Then in $\triangle GCD$ are given GC, CD and $\angle CGD$ to get GD. And now GA and GB follow in a very obvious manner.

This problem may obviously be solved with fewer equations by using quadrilaterals, instead of triangles, in the solution.

Unless the two circles of position are tangent to each other there will be two points (viz, G and G') which will equally well satisfy the conditions of the problem: hence in this case we judge from an approximate knowledge of our position which of the points G or G' we were at when we observed.

The manner of sweeping the circles of position, with Protractor and Dividers is easy and expeditious: since $\angle AEB = 2 \angle AGB$ then $\angle ABE$

$$= \frac{180^\circ - \angle AEB}{2} = \frac{180^\circ - 2 \angle AGB}{2} = 90^\circ - \angle AGB,$$

that is $\angle ABE =$ the complement of the observed $\angle AGB$. The rule for plotting the circles of position is now obviously, to lay off from AB at the points A, and B the complement of the observed angle AGB, and the point of intersection of the produced sides of these angles will be

the center of a circle of position. And now from this point as center with radius AE (or BE) sweep the circle of position. And in like manner lay down the other circle of position through C and D.

This problem will often be found servicable to the Hydrographer and Explorer when from either accident or necessity only two angles are measured on four objects.

SOLUTIONS OF PROBLEMS IN NO. 6.

Solutions of problems in No. 6 have been received as follows: From Geo. L. Dake, 25 & 26; R. M. DeFrance, 25 & 26; Prof. A. B. Evans, 25, 26, 27 & 29; Henry Gunder, 25, 26, 27 & 29; Wm. Hoover, 26; Prof. A. Hall, 29; H. Heaton, 29; D. J. McAdam, 25 & 26; Esther Matthews, 26; Artemas Martin, 27 & 29; A. W. Phillips, 25, 26, 27 & 29; L. Regan, 25, 26 & 29; R. L. Selden, 25; Werner Stille, 25, 26, 27 & 29; E. B. Seitz, 25, 26, 27 & 29; Prof. J. Scheffer, 26 & 29; Walter Siverly, 27.

25. "Required the sides of an obtuse angled triangle the area of which is 14.048 acres, the obtuse angle $111^{\circ}15'$, and one of the acute angles $11^{\circ}44'10''$."

SOLUTION BY HENRY GUNDER, GREENVILLE, OHIO.

Putting $A = 111^{\circ}15'$, $B = 11^{\circ}44'10''$, $C = 57^{\circ}50''$, and x, y, z or the sides opposite A, B and C , and $a = 14.048$ acres $= 2247.68$ sq. rods.

Since the product of two sides and the sine of the included angle equals twice the area we have,

$$(1) xy = \frac{2a}{\sin C}, \quad (2) xz = \frac{2a}{\sin B}, \quad (3) yz = \frac{2a}{\sin A}. \quad \text{Then}$$

$$\sqrt{\frac{(1) \times (2)}{(3)}} = (4) x = \sqrt{\frac{2a \sin A}{\sin B \sin C}}. \quad \text{Similarly } y = \sqrt{\frac{2a \sin B}{\sin A \sin C}}.$$

$$\text{" } z = \sqrt{\frac{2a \sin C}{\sin A \sin B}}.$$

By applying logarithms, $x = 156.705$ rods, $y = 34.1997$ rods, $z = 141.034$ rods.